

### **Drug Description: TADACIP**

Due to its quick and spontaneous action and long-lasting effects, today Cialis is considered to be one of the most popular erectile dysfunction drugs in the world. For the last few years, numerous clinical trials have already been conducted with the primary aim of developing generic versions of Cialis (Tadalafil). Being one of the most reputed manufacturers of generic medication, Cipla developed the Tadacip (Tadalafil), which has become the topmost erectile dysfunction drug among the other generics range of medications. People who have already used it have confirmed about the effectiveness and power of Tadacip (Tadalafil). The phosphodiesterase type 5(PDE-5) inhibitor happens to be the most important component for most of these erectile dysfunction drugs. The erectile dysfunction (ED), also called as the male impotence is characterized by the inability of a person to achieve erection or maintain erection until climax during the sexual intercourse. ED can also lead to other diseases such as high blood pressure, diabetes, vascular disease, nervous system disorder, heart disease and depression. This erectile dysfunction drug works better than the other Cialis medications and also that too for half the price of the other Cialis medications.

**Availability:** In stock

**Manufacturer:** Cipla

**Sold Under Brand Name:** Tadacip, Cialis

**Average Delivery Time:** 8 days

#### **Benefits**

- Tadacip is a much better option than the Generic Viagra
- People under the medication of Tadacip do not have to go through the pain of planning their sexual intercourse
- The effects of Tadacip (Tadalafil) lasts for a longer period unlike the other erectile dysfunction drugs often found in the market or in the online pharmacy web portals. The effects stays for as long as 36 hours

- Fatty foods do not dispel the effects of Tadacip, so it can be taken with or without a meal
- Tadacip starts working just after half an hour after consumption

## **Product Information**

### **How to use**

One can take Tadacip (Tadalafil) anytime in 24 hours for once or he can follow the directions of the physician. Tadacip's each dose should be taken orally along with a full glass of water half an hour before the sexual intercourse. This erectile dysfunction drug can be taken with or without food. Tadacip works effectively only when sexual stimulation occurs. Don't expect the medication to work without any kind of sexual urge.

### **Overdose**

Dizziness, irregular heartbeat, light-headedness and nausea are some of the common symptoms of Tadacip (Tadalafil) overdose. One should immediately get in touch with the nearest poison control center or contact the emergency medical services in order to deal with the situation immediately.

### **Missed dose**

Tadacip has to be consumed just before sexual intercourse. Thus, there is a frail chance of missing its dose.

### **Notes**

Tadacip (Tadalafil) cannot be shared with anyone else who does not have this particular malfunction. Tadacip also does not help in preventing the spread of the sexually transmitted diseases like gonorrhea, syphilis etc.

### **Storage**

- Tadacip (Tadalafil) should always be kept out of reach from the children
- Cialis Super Active (Tadalafil) should be stored in a room with a temperature between 59 and 86 degrees F (15-30 degrees C).
- Tadacip (Tadalafil) should always be stored away from light and moisture

### **Important note**

A physician, pharmacist or any other healthcare professional's directions are the most important ones. The information provided above intends to aid in the knowledge of the people Tadacip (Tadalafil).

This document is protected by copyright in the name of the company. It is not meant for re-use in any form and by any means.

## **FAQs**

### **Is alcohol consumption safe while under the medication of Tadacip (Tadalafil)?**

Excessive alcohol consumption while under the medication of Tadacip can increase the risk of certain side effects, such as headache, dizziness, increased heart rate, or low blood pressure. Thus, alcohol consumption is not at all a good idea while taking Tadacip.

### **What happens if I miss a dose?**

There is no question of missing the dose as one can automatically remember the need of taking the medicine just before sexual intercourse.

### **How does Tadacip (Tadalafil) work?**

Due to the presence of the phosphodiesterase inhibitors, Tadacip helps in relaxing the muscles and increases blood flow to the penis muscles. This helps in achieving and maintaining an erection for a longer time.

## **Precautions**

Do inform your doctor if you have the given below malfunctions in your body before including Tadacip (Tadalafil) in your medication routine –

- Heart problems such as angina, heart failure or irregular heartbeat or have had a heart attack
- Recent (last 90 days) report of a heart attack
- A report of stroke or congestive heart failure in the past 6 months
- Blood cell problems such as leukemia, multiple myeloma or sickle cell anemia
- Low blood pressure or high blood pressure that is not controlled
- Bleeding disorders such as hemophilia
- Liver or kidney problems
- A stomach ulcer
- Retinitis pigmentosa (an condition of the eye the patient usually inherits)
- A physical defect of the penis such as Peyronie's disease
- Erection that lasted longer than 4 hours
- Never take this medicine if you are asked for not taking due health reasons
- Do not use this medication without informing your doctor if you are breast-feeding.

This document is protected by copyright in the name of the company. It is not meant for re-use in any form and by any means.

## **Drug Interactions**

If you are the given below medications, do not prescribe to this medicine –

- Alpha blockers (other than tamsulosin (Flomax) 0.4 mg once a day) such as doxazosin (Cardura), guanadrel (Hylorel), prazosin (Minipress), terazosin (Hytrin), alfuzosin (Uroxatral) and others.
- Nitrates found in some recreational drugs such as amyl nitrate or nitrite ("poppers")
- Nitrates such as nitroglycerin (Nitrostat, Nitrolingual, Nitro-Dur, Nitro-Bid, Minitran, Deponit, Transderm-Nitro, others), isosorbide dinitrate (Dilatrate-SR, Isordil, Sorbitrate), isosorbide mononitrate (Imdur, ISMO, Monoket) and others
- Antifungal medications such as clotrimazole (Mycelex Troche), itraconazole (Sporanox), ketoconazole (Nizoral) or voriconazole (Vfend)
- Antidepressants such as nefazodone
- Conivaptan (Vaprisol)
- Seizure medications such as carbamazepine (Tegretol), phenobarbital (Luminal), or phenytoin (Dilantin)
- Diclofenac (Arthrotec, Cataflam, Voltaren, Flector Patch, Solareze)
- HIV/AIDS medications, such as amprenavir (Agenerase), atazanavir (Reyataz), darunavir (Prezista), delavirdine (Rescriptor), fosamprenavir (Lexiva), indinavir (Crixivan), nelfinavir (Viracept), saquinavir (Invirase, Fortovase) or ritonavir (Norvir)
- Doxazosin (Cardura), prazosin (Minipress), terazosin (Hytrin)
- Heart or blood pressure medications
- Imatinib (Gleevec)
- Isoniazid (for treating tuberculosis)
- Rifampin (Rifadin, Rimactane)
- An antibiotic such as clarithromycin (Biaxin), dalfopristin/quinupristin (Synercid), erythromycin (E.E.S., EryPed, Ery-Tab, Erythrocin) or telithromycin (Ketek)

## **Side-Effects**

The following are some of the common side-effects that may show up if someone is under the medication of Tadacip (Tadalafil) –

- Shortness of breath
- Dizziness, nausea, or angina (pain, tightness, discomfort, numbness, or tingling in the chest, arms, neck or jaw)

This document is protected by copyright in the name of the company. It is not meant for re-use in any form and by any means.

- Allergic reactions (difficulty breathing, closing of the throat, swelling of the lips, tongue or face, hives)
- Prolonged or severe dizziness
- Swelling of the ankles or legs
- Chest pain or irregular heartbeat
- A prolonged (lasting longer than 4 hours) or painful erection

There are some other unlikely side-effects also –

- Temporary blue tint in vision or difficulty telling the difference between the colors blue and green (uncommon)
- Flushing
- Back pain
- Muscle aches
- Indigestion
- Headache
- Stuffy or runny nose

In both the cases, a physician or a healthcare professional should be immediately contacted to avoid further complications.

### **Warnings**

- The side effects cited above for Tadacip (Tadalafil) may not cover all possible side effects.
- The symptom of overdose of Tadacip depends upon the reaction of the body to the erectile dysfunction drug.
  - In case, if the side effects of Tadacip worsen, one needs to contact the doctor or medical emergency unit immediately.